



BEEF COW PROTOCOLS - 2017

HEAT DETECTION

Select Synch




Select Synch + CIDR®



PG 6-day CIDR®


Heat detect and AI days 0 to 3. Administer CIDR to non-responders and heat detect and AI days 9 to 12. Protocol may be used in heifers.



HEAT DETECT & TIME AI (TAI)


Select Synch & TAI

Heat detect and AI day 6 to 10 and TAI all non-responders 72 - 84 hr after PG with GnRH at TAI.




Select Synch + CIDR® & TAI

Heat detect and AI day 7 to 10 and TAI all non-responders 72 - 84 hr after PG with GnRH at TAI.



PG 6-day CIDR® & TAI


Heat detect & AI days 0 to 3. Administer CIDR to non-responders & heat detect and AI days 9 to 12. TAI non-responders 72 - 84 hr after CIDR removal with GnRH at AI. Protocol may be used in heifers.



FIXED-TIME AI (TAI)*


7-day CO-Synch + CIDR®

Perform TAI at 60 to 66 hr after PG with GnRH at TAI.



5-day CO-Synch + CIDR®

Perform TAI at 72 ± 2 hr after CIDR removal with GnRH at TAI. Two injections of PG 8 ± 2 hr apart are required for this protocol.




FIXED-TIME AI (TAI)*

for *Bos Indicus* cows only

PG 5-day CO-Synch + CIDR®

Perform TAI at 66 ± 2 hr after CIDR removal with GnRH at TAI. Two injections of PG 8 ± 2 hr apart are required for this protocol.




* The time listed for "Fixed-time AI" should be considered as the approximate average time of insemination. This should be based on the number of cows to inseminate, labor, and facilities.

GnRH Cystorelin®, Factrel®, Fertagyl®, OvaCyst®, GONABreed®
PG estroPLAN®, Estrumate®, In-Synch®, Lutalyse®, Lutalyse® HighCon, ProstaMate®, SYNCHSURE™


BEEF HEIFER PROTOCOLS - 2017

HEAT DETECTION


1 Shot PG



7-day CIDR®-PG




MGA®-PG



HEAT DETECT & TIME AI (TAI)


Select Synch + CIDR® & TAI

Heat detect and AI day 7 to 10 and TAI all non-responders
72 - 84 hr after PG with GnRH at TAI.




MGA®-PG & TAI

Heat detect and AI day 33 to 36 and TAI all non-responders
72 - 84 hrs after PG with GnRH at TAI.



14-day CIDR®-PG & TAI

Heat detect and AI day 30 to 33 and TAI all non-responders
72 hrs after PG with GnRH at TAI.




FIXED-TIME AI (TAI)*

Short-term Protocols


7-day CO-Synch + CIDR®

Perform TAI at 54 ± 2 hr after PG with GnRH at TAI.



5-day CO-Synch + CIDR®


Perform TAI at 60 ± 4 hr after CIDR removal with GnRH at TAI.
Two injections of PG 8 ± 2 hr apart are required for this protocol.



Long-term Protocols


14-day CIDR®-PG

Perform TAI at 66 ± 2 hr after PG with GnRH at TAI.




MGA®-PG

Perform TAI at 72 ± 2 hr after PG with GnRH at TAI.



* The times listed for "Fixed-time AI" should be considered as the approximate average time of insemination. This should be based on the number of heifers to inseminate, labor, and facilities.

 Cystorelin®, Factrel®, Fertagyl®, OvaCyst®, GONABreed®

 estroPLAN®, Estrumate®, In-Synch®, Lutalyse®, Lutalyse® HighCon, ProstaMate®, SYNCHSURE™